

Radiative stability of neutrino-mass textures

M. K. Parida* and C. R. Das†

Department of Physics, North Eastern Hill University, Shillong 793022, India

G. Rajasekaran‡

Institute of Mathematical Sciences, Chennai 600113, India

(Dated: February 1, 2008)

Neutrino-mass textures proposed at high-scales are known to be unstable against radiative corrections especially for nearly degenerate eigen values. Within the renormalization group constraints we find a mechanism in a class of gauge theories which guarantees reproduction of any high-scale texture at low energies with radiative stability. We also show how the mechanism explains solar and atmospheric neutrino anomalies through the bimaximal texture at high scale.

PACS numbers: 14.60.Pq, 11.30.Hv, 12.15.Lk

I. Introduction: A major challenge to particle physics at present is the theoretical understanding of experimentally observed neutrino anomalies. This has led to suggestions of many interesting models and mass textures with hierarchical or degenerate eigen values [1, 2, 3, 4, 5, 6, 7, 8]. Whereas the observed mixing between quarks are small, experimental indications appear to favor maximal mixings in the neutrino sector. A possible mechanism to explain large neutrino mixings at low-energies starting from small high-scale mixings similar to quarks could be through radiative magnification for quasidegenerate neutrinos [9, 10].

An outstanding problem with bimaximal neutrino-mass textures with degenerate eigen values is the instability of the masses and mixing angles due to radiative corrections which spoils their prospects for the neutrinoless-double-beta decay and the neutrino anomalies [10, 11, 12]. While investigating radiative stability the usual procedure has been to assume the bimaximal texture to be associated with a single 5-dim operator through see-saw mechanism with SM or MSSM as gauge theories at lower scales [9, 10, 11, 12].

Using renormalization group constraints in this letter we show that it is possible to reproduce any high-scale mass texture at low scale ($\mu = M_Z$) with high degree of accuracy leading to stable evolution of the physically relevant Majorana-neutrino-mass matrix. The models where the mechanism operates consists of two component matrices contributing to the resultant Majorana-neutrino

mass texture at the highest scale. It is quite interesting to note that the mechanism operates successfully in 2HDM and also with SM and MSSM in the presence of type II see-saw mechanism probing left-right model (LRM) and $SO(10)$ GUT as prospective high-scale theories [6, 13, 14]. We also show how experimental data on solar and atmospheric neutrino anomalies are explained through the bimaximal texture.

II. The mechanism in 2HDM: In the SM, MSSM, and 2HDM, the Majorana-neutrino mass in the flavor basis originates from 5-dim operators generated at the lepton-number-breaking scale ($\mu_0 \simeq M_N$, $t_0 = \ln M_N$) through see-saw mechanism. In the SM there is only one Higgs doublet Φ with VEV $\langle \Phi \rangle_0 = v/\sqrt{2}$ and one operator ($= K^{(SM)}$) contributing to the Majorana mass $m^{(SM)} = -(1/4)K^{(SM)}v^2$. In the MSSM, although there are two Higgs doublets, there is only one neutrino-mass operator contributing to the neutrino-mass matrix due to the coupling of up type doublet with $m^{(MSSM)} = -(1/4)K^{(MSSM)}v_u^2$. In a class of 2HDM there are two doublets, Φ_u and Φ_d , with VEVs $v_u/\sqrt{2} = v \sin \beta/\sqrt{2}$, $v_d/\sqrt{2} = v \cos \beta/\sqrt{2}$. But, unlike SM or MSSM, there are two neutrino-mass operators, K^I and K^{II} , and two matrices m^I and m^{II} which add up to generate the physically relevant Majorana-neutrino-mass matrix [10], $m = -(1/4)(K^I v_u^2 + K^{II} v_d^2) \equiv m^I + m^{II}$. We use the renormalization scheme where the runnings of v_u , v_d , and $\tan \beta$ are ignored. Then the relevant RGEs and their one-loop solutions for $\mu < M_N$ ($t = \ln \mu < t_0$) are,

$$16\pi^2 \frac{dm^I}{dt} = \left\{ -3g_2^2 + 2\lambda_2 + 2\text{Tr} \left(3Y_U^\dagger Y_U \right) \right\} m^I + \frac{1}{2} \left[\left(Y_E^\dagger Y_E \right) m^I + m^I \left(Y_E^\dagger Y_E \right)^T \right] + 2\lambda_3 m^{II},$$

$$16\pi^2 \frac{dm^{II}}{dt} = \left\{ -3g_2^2 + 2\lambda_1 + 2\text{Tr} \left(3Y_D^\dagger Y_D + Y_E^\dagger Y_E \right) \right\} m^{II} - \frac{3}{2} \left[\left(Y_E^\dagger Y_E \right) m^{II} + m^{II} \left(Y_E^\dagger Y_E \right)^T \right] + 2\lambda_3^* m^I,$$

$$m_{ij}^I(t) = a_{ij}^I(t) m_{ij}^I(0), \quad a_{ij}^I(t) = I_{g_2}^{-\frac{3}{2}} I_{\lambda_2} I_{\tan \beta}^3 (I_i I_j)^{\frac{1}{4}} R_{ij}, \quad (1)$$

$$m_{ij}^{II}(t) = a_{ij}^{II}(t) m_{ij}^{II}(0), \quad a_{ij}^{II}(t) = I_{g_2}^{-\frac{3}{2}} I_{\lambda_1} I_b^3 I_\tau (I_i I_j)^{-\frac{3}{4}} \tilde{R}_{ij}. \quad (2)$$

Here $m_{ij}^{I,II}(0) = m_{ij}^{I,II}(t_0)$,

$$\begin{aligned} I_l &= \exp\left(\frac{1}{8\pi^2} \int_{t_0}^t h_l^2 dt\right), (l = e, \mu, \tau, \text{top}, b) \\ I_{g_k} &= \exp\left(\frac{1}{8\pi^2} \int_{t_0}^t g_k^2 dt\right), (k = 1, 2) \\ I_{\lambda_k} &= \exp\left(\frac{1}{8\pi^2} \int_{t_0}^t \lambda_k dt\right), (k = 1, 2) \\ R_{ij} &= \exp\left[\frac{1}{8\pi^2} \int_{t_0}^t \left(m^{II} m^{I-1}\right)_{ij} \lambda_3 dt\right], \\ \tilde{R}_{ij} &= \exp\left[\frac{1}{8\pi^2} \int_{t_0}^t \left(m^I m^{II-1}\right)_{ij} \lambda_3^* dt\right]. \end{aligned} \quad (3)$$

Any texture at the highest scale for the physically relevant Majorana-neutrino-mass matrix

$$m(0) = m^I(0) + m^{II}(0), \quad (4)$$

never determines both the matrices $m^I(0)$ and $m^{II}(0)$. Given any element $m_{ij}(0)$ at t_0 , one of the component elements, $m_{ij}^I(0)$ or $m_{ij}^{II}(0)$, remains completely undetermined at that scale. Then (1)-(2) show that the same matrix, $m^I(t)$ or $m^{II}(t)$, is undetermined at all lower scales too. This is in clear contrast to the cases in conventional analyses (CA) with SM or MSSM where there is only one $m(0)$ at $\mu = M_N$ and the texture gives all the elements of $m_{ij}(0)$ and $m_{ij}(t)$ [10, 11]. Now we impose the stability criterion that the texture is exactly reproduced at the lowest scale by demanding that

$$m_{ij}(t_Z) = m_{ij}^I(t_Z) + m_{ij}^{II}(t_Z) \equiv m_{ij}(0). \quad (5)$$

Since $a^I(t_Z)$ and $a^{II}(t_Z)$ are known in terms of the model parameters, solutions of (4) and (5) now determine both $m^I(0)$ and $m^{II}(0)$ in terms of the high-scale neutrino-mass texture, $m(0)$,

$$\begin{aligned} m_{ij}^I(0) &= m_{ij}(0) (a_{ij}^{II}(t_Z) - 1) / d_{ij}, \\ m_{ij}^{II}(0) &= m_{ij}(0) (1 - a_{ij}^I(t_Z)) / d_{ij}, \\ d_{ij} &= a_{ij}^{II}(t_Z) - a_{ij}^I(t_Z). \end{aligned} \quad (6)$$

These parameters of the component matrices, determined from the boundary conditions (4) and (5) are expected to guarantee reproduction of the high-scale texture at M_Z when $m^I(t)$ and $m^{II}(t)$ are evolved through (1)-(3).

As an example we study RG evolution of the bimaximal texture with triply degenerate masses at $M_N \simeq 10^{13}$ GeV [4]

$$m(0) = \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{\sqrt{2}} & -\frac{1}{2} & \frac{1}{2} \end{bmatrix} m_0. \quad (7)$$

Using, $\lambda_1 = 0.16$, $\lambda_2 = 1.13$, $\lambda_3 \simeq -0.011$ [10], and $\tan \beta = 40$ we obtain $I_{\lambda_1} \simeq 0.95$, $I_{\lambda_2} \simeq 0.6976$, $I_{\text{top}} =$

0.833213, $I_b = 0.935023$, $I_\tau = 0.950882$, $I_\mu = 0.999832$, $I_e = 0.999999$, $I_{g_2} = 0.478614$ at $\mu = M_Z$. We compute $a^I(t_Z)$ and $a^{II}(t_Z)$, and, hence, $m^I(0)$ and $m^{II}(0)$ shown in Table I as input parameters. The solutions for $m^I(t)$ and $m^{II}(t)$ are obtained through (1)-(3) and the elements of the Majorana-neutrino-mass matrix $m_{ij}(t)$ are obtained as their sum for all $t < t_0$. In Fig. 1 we have shown the radiative corrections for $m_{\tau\tau}(t)$. For comparison we have shown the results of the conventional analysis as SM (CA) and MSSM (CA) for which there is only one matrix at the highest scale. The maximum radiative correction of the matrix elements in 2HDM using the present mechanism is found to be only 3-4% as compared to 30-40% in the SM (CA) or MSSM (CA). Whereas the maximal corrections in SM (CA) or MSSM (CA) occur at $\mu = M_Z$, in our case they occur with substantially reduced magnitude at intermediate scales. NonSUSY SM and 2HDM have been successfully embedded in $SO(10)$ with single intermediate symmetries [15].

III. Implementation in SM or MSSM: We note that the present mechanism also operates in SM and MSSM if they originate from high-scale theories which predict two component matrices at M_N . The popular see-saw mechanism which has its natural origin in LRM and $SO(10)$ contains the second contribution and leads to the two matrices in type II see-saw formula with $m^I = m^{\text{SM}}(m^{\text{MSSM}})$ and $m^{II} \simeq f v^2 / M_N (f v_u^2 / M_N)$ when SM (MSSM) is obtained after symmetry breaking of LRM or $SO(10)$ [1, 6, 7, 8, 14, 15, 16]. Here f is the Majorana type Yukawa coupling of the neutrino. The mechanism also operates in SM or MSSM when there are other types of contributions [16]. The problem of obtaining a specific texture for $m(0)$, or $m^I(0)$ and $m^{II}(0)$, may call for appending specific flavor symmetries to $SU(2)_L \times SU(2)_R \times U(1)_{B-L} \times SU(3)_C$ or $SO(10)$. Assuming such possibilities we derive the constraints on $m^I(0)$ and $m^{II}(0)$ resulting from the bimaximal texture for $m(0)$ and its radiative stability. The RG evolutions of the standard see-saw term is the same as in SM or MSSM as shown through $a^I(t)$ in (8)-(9) below. But those for $m^{II}(t)$ occur due to loop-mediation of the standard-weak-Higgs doublet and gauge bosons (plus superpartners) with the LH neutrinos alone. We derive them as

$$\begin{aligned} 16\pi^2 \frac{dm^{II}}{dt} &= \left(c^{(1)} g_1^2 + c^{(2)} g_2^2\right) m^{II} \\ &+ c^{(3)} \left[\left(Y_E^\dagger Y_E\right) m^{II} + m^{II} \left(Y_E^\dagger Y_E\right)^T \right], \end{aligned}$$

where $c^{(i)} = (9/10, 3/2, -3/2)$ for SM, but $c^{(i)} = (-9/5, -9/2, 1)$ for MSSM. In the notations of (1)-(2) we obtain

SM

$$a_{ij}^I(t) = I_{g_2}^{-\frac{3}{2}} I_{\lambda_{\text{top}}}^3 I_b^3 I_\tau (I_i I_j)^{-\frac{3}{2}},$$

$$a_{ij}^{II}(t) = I_{g1}^{\frac{9}{20}} I_{g2}^{\frac{3}{4}} (I_i I_j)^{-\frac{3}{4}}, \quad (8)$$

MSSM

$$\begin{aligned} a_{ij}^I(t) &= I_{g1}^{-\frac{3}{5}} I_{g2}^{-3} I_{top}^3 (I_i I_j)^{\frac{1}{2}}, \\ a_{ij}^{II}(t) &= I_{g1}^{-\frac{9}{10}} I_{g2}^{-\frac{9}{4}} (I_i I_j)^{\frac{1}{2}}. \end{aligned} \quad (9)$$

Then using (4)-(6) we obtain the initial values of $m^{I,II}(0)$ and, hence, solutions for $m_{ij}(t)$ for $\mu = M_Z - M_N$ exhibiting stability of all the matrix elements of $m(t)$ under radiative corrections. The elements of the component matrices for the two cases are also shown in Table I. In Fig. 1 we have plotted $m_{\tau\tau}(t)$ in comparison to conventional analyses. As against the maximal 30-40% radiative corrections in SM (CA) and MSSM (CA) occurring at $\mu = M_Z$, they are only 3-4% in SM and 1.5% in MSSM which occur at intermediate scales in the present analysis. Among all the three models, the minimum radiative corrections upto 1.5% is found to occur in MSSM.

IV. Fitting the neutrino anomalies: When the bi-maximal texture is exactly reproduced at M_Z , one way to explain neutrino anomalies could be through threshold effects [17]. But here ignoring threshold effects we show how the present mechanism permits matching of the observed solar (LAMSW) and atmospheric neutrino anomalies starting from the bi-maximal texture with degenerate mass eigen values at the highest scale. Using quasidegenerate neutrinos with masses $m_1 = -0.2$ eV, $m_2 = 0.200045$ eV, $m_3 = 0.2075$ eV which are spread around $m_0 = 0.2$ eV, the mixing angles suitable for LAMSW with $s_3 = 0.6946$ and atmospheric neutrino oscillations with $s_1 = 0.6950$, it is straight forward to construct the mass matrix consistent with the experimental data

$$m^{(e)}(t_Z) = \begin{bmatrix} -0.044716 & 0.722025 & -0.690426 \\ 0.722025 & 0.501718 & 0.477908 \\ -0.690426 & 0.477908 & 0.544506 \end{bmatrix} m_0. \quad (10)$$

Although we have used $s_2 = 0$, the mechanism is found to work for other values consistent with CHOOZ bound [18]. Similarly the mechanism also works with other values of $m_0 \simeq 0.1 - 1.0$ eV. Within the RG-constraints, the high scale texture can match the experimentally observed anomalies provided $m_{ij}(0)$ in (5) and (6) is replaced by $m_{ij}^{(e)}(t_Z)$ leading to

$$\begin{aligned} m_{ij}^I(0) &= \left(a_{ij}^{II}(t_Z) m_{ij}(0) - m_{ij}^{(e)}(t_Z) \right) / d_{ij}, \\ m_{ij}^{II}(0) &= \left(m_{ij}^{(e)}(t_Z) - a_{ij}^I(t_Z) m_{ij}(0) \right) / d_{ij}. \end{aligned} \quad (11)$$

In Fig. 1 the curves 2HDM(e), MSSM(e) and SM(e) represent the result of fitting the data through the high scale bi-maximal texture given in (7) and $m^{(e)}(t_Z)$ given in (10) using 2HDM, MSSM and SM, respectively. We note that similar RG-stability also holds approximately for certain




FIG. 1: RG-stable evolutions of $m_{\tau\tau}(t)$ in 2HDM, MSSM with $\tan\beta = 40$, and SM. MSSM (CA) and SM (CA) denote conventional analyses in MSSM and SM having only one matrix at the highest scale. 2HDM (e), MSSM (e) and SM (e) represent RG evolutions matching the experimental data on neutrino anomalies.

other elements depending upon the exact values of s_1 and s_3 . But the radiative corrections are found to be larger if the difference between s_1 and s_3 is larger. Similar curves can be plotted for other elements also.

V. Conclusion: The present mechanism demonstrates how to evade RG-constraints on neutrino-mass textures in conventional analyses. It operates in a class of gauge theories leading to 2HDM, SM or MSSM where two component matrices contribute to the physically relevant Majorana-neutrino mass at the highest scale. Once a resultant texture is generated using suitable flavor symmetries at the highest scale, this mechanism determines the two unknown matrices at the highest scale which ensure its RG-stability at all lower scales or its matching with the experimental data. The mechanism can be applied to reproduce any high-scale texture at low energies with any desired degree of stability including higher order corrections in (5). It is quite interesting that the stability criteria operate in the presence of type II see-saw mechanism and probe into models including left-right gauge theories and $SO(10)$ as prospective high-scale theories. The textures for component matrices derived from the stability condition sets considerable constraint on future model building with flavor symmetry.

The work of M.K.P. is supported by Project No. SP/S2/K-30/98 and the work of C.R.D. is supported by Project No. 98/37/9/BRNS-Cell/731 of the Government of India.

* Electronic address: mparida@nehu.ac.in; Electronic address: mparida@sancharnet.in

† Electronic address: crdas@email.com

‡ Electronic address: graj@imsc.ernet.in

[1] R.N. Mohapatra, hep-ph/9910365, hep-ph/0008232; S.

TABLE I: Component matrices determined from RG stability criteria and also by matching the experimental data as denoted by (e). Here m_0 is a common factor.

Model	$m^I(0)/m_0$	$m^{II}(0)/m_0$
2HDM	$\begin{bmatrix} 0.000000 & -0.220454 & -0.146400 \\ -0.220454 & -0.155695 & 0.103358 \\ -0.146400 & 0.103358 & -0.058182 \end{bmatrix}$	$\begin{bmatrix} 0.000000 & 0.927561 & 0.853507 \\ 0.927561 & 0.655695 & -0.603358 \\ 0.853507 & -0.603358 & 0.558182 \end{bmatrix}$
2HDM (e)	$\begin{bmatrix} 0.112140 & -0.257854 & 3.041252 \\ -0.257854 & -0.160003 & -2.126483 \\ 3.041252 & -2.126483 & -0.151038 \end{bmatrix}$	$\begin{bmatrix} -0.112140 & 0.964961 & -2.334145 \\ 0.964961 & 0.660003 & 1.626483 \\ -2.334145 & 1.626483 & 0.651038 \end{bmatrix}$
MSSM	$\begin{bmatrix} 0.000000 & 0.414613 & 0.389877 \\ 0.414613 & 0.293120 & -0.275628 \\ 0.389877 & -0.275628 & 0.257691 \end{bmatrix}$	$\begin{bmatrix} 0.000000 & 0.292493 & 0.317228 \\ 0.292493 & 0.206879 & -0.224371 \\ 0.317228 & -0.224371 & 0.242308 \end{bmatrix}$
MSSM (e)	$\begin{bmatrix} 0.061597 & 0.394061 & 2.364023 \\ 0.394061 & 0.290752 & -1.657126 \\ 2.364023 & -1.657126 & 0.193220 \end{bmatrix}$	$\begin{bmatrix} -0.061597 & 0.313044 & -1.656916 \\ 0.313044 & 0.209247 & 1.157126 \\ -1.656916 & 1.157126 & 0.306779 \end{bmatrix}$
SM	$\begin{bmatrix} 0.000000 & -0.292357 & -0.292451 \\ -0.292357 & -0.206728 & 0.206794 \\ -0.292451 & 0.206794 & -0.206860 \end{bmatrix}$	$\begin{bmatrix} 0.000000 & 0.999463 & 0.999557 \\ 0.999463 & 0.706728 & -0.706794 \\ 0.999557 & -0.706794 & 0.706860 \end{bmatrix}$
SM (e)	$\begin{bmatrix} 0.162944 & -0.346719 & 4.800250 \\ -0.346719 & -0.212991 & -3.356768 \\ 4.800250 & -3.356768 & -0.369052 \end{bmatrix}$	$\begin{bmatrix} -0.162944 & 1.053825 & -4.093143 \\ 1.053825 & 0.712991 & 2.856768 \\ -4.093143 & 2.856768 & 0.869052 \end{bmatrix}$

- Barr and I. Dorsner, hep-ph/0003058.
- [2] R. Barbieri et. al., hep-ph/9901228, hep-ph/9807235, hep-ph/9906470; N. Irges, S. Lavignac, and P. Ramond, hep-ph/9802334.
- [3] K.S. Babu, J.C. Pati and F. Wilczek, hep-ph/9812538.
- [4] H. Georgi and S.L. Glashow, hep-ph/9808293; V. Barger et. al., hep-ph/9806387; G. Altarelli and F. Feruglio, hep-ph/9807353; R.N. Mohapatra and S. Nussinov, hep-ph/9808301; F. Vissani, hep-ph/9708483.
- [5] C.H. Albright and S.M. Barr, hep-ph/9712488, hep-ph/0002155, hep-ph/0003251; C.H. Albright, K.S. Babu, and S.M. Barr, hep-ph/9802314; T. Blazek, S. Raby and K. Tobe, hep-ph/9903340; M. Chen and K.T. Mahanthappa, hep-ph/0009059.
- [6] K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **70**, 2845 (1993); B. Brahmachari and R.N. Mohapatra, Phys. Rev. D **58**, 015001 (1998); C.S. Aulakh, B. Bajc, A. Melfo, A. Rasin and G. Senjanovic, hep-ph/0004031.
- [7] D. Caldwell and R.N. Mohapatra, Phys. Rev. D **48**, 3259 (1993), Phys. Rev. D **50**, 3477 (1994).
- [8] A. Ioannissyan and J. Valle, Phys. Lett. B **332**, 93 (1994); J. Peltoniemi and J.W.F. Valle, Nucl. Phys. B **406**, 409 (1993).
- [9] K.R.S. Balaji, A.S. Dighe, R.N. Mohapatra, and M.K. Parida, Phys. Rev. Lett. **84**, 5034 (2000); Phys. Lett. B **481**, 33 (2000); K.R.S. Balaji, R.N. Mohapatra, M.K. Parida, and E.A. Paschos, Phys. Rev. D **63**, 113002 (2001).
- [10] K.S. Babu, C.N. Leung, and J. Pantaleone, Phys. Lett. B **319**, 191 (1993); S. Antusch, et. al, hep-ph/0108005; hep-ph/0110366.
- [11] J.Ellis and S. Lola, hep-ph/9904279; J.A. Casas et. al. hep-ph/9904395, hep-ph/9905381, hep-ph/9906281, hep-ph/9910420; P.H. Chankowski et. al., hep-ph/9910231; N. Haba et. al., hep-ph/9810471, hep-ph/9905381; P.H. Chankowski and Z. Pluciennik, Phys. Lett. B **316**, 312 (1993); M.K. Parida and N.N. Singh, Phys. Rev. D **59**, 032002 (1999).
- [12] E. Ma, J. Phys. G **25**, L97 (1999); R. Adhikari, E. Ma, and G. Rajasekaran, hep-ph/0004197.
- [13] M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, eds. P. Van Nieuwenhuizen and D.Z. Freedman (North-Holland, Amsterdam, 1979) p. 315; T. Yanagida, Proc. Workshop on Unified theory and baryon number of the universe (Tsukuba, Ibaraki, Japan), unpublished (1979), R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [14] R.N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981).
- [15] D. Chang, R.N. Mohapatra, and M.K. Parida, Phys. Rev. Lett. **52**, 1072 (1984); Dae-Gye Lee, R.N. Mohapatra, M.K. Parida, and M. Rani, Phys. Rev. D **51**, 229 (1995); M.K. Parida and A. Usmani, Phys. Rev. D **54**, 3663 (1996).
- [16] E. Ma and U. Sarkar, Phys. Rev. Lett. **80**, 5716 (1998).
- [17] P.H. Chankowski et. al., hep-ph/0011150; P.H. Chankowski and P. Wasowicz, hep-ph/0110237.
- [18] M. Apollonio et. al., hep-ex/9907037.